


MAR 06 2008

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently Amended) A compound of formula I



formula I

Wherein, n is 0 or 1;

R1 is XR wherein;

X is selected from O; S; S=O and SO₂,

R is a moiety selected from the group consisting of: C1 to C9 alkyl, C2 to C15 alkenyl, C2 to C15 alkynyl, C1 to C15 heteroalkyl, C6 to C15 aryl, C6 to C15 heteroaryl, C6 to C15 arylalkyl or C6 to C15 heteroarylalkyl; which moiety R is optionally substituted, cyclic or acyclic, branched and/or linear,

the groups R2 to R5 are selected from OH, OR and N(Y)Z such that:


at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OH,

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OR, where R is defined above, with the proviso that when two of the

groups R2 to R5 are OR, OR is an ether type moiety and the R groups may not both be methyl, C1 to C3 n-alkyl, allyl or unsubstituted benzyl,

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;

wherein when n=1, at least one of the groups R4 or R5 is OR or N(Y)Z;



WEST et al
Appl. No. 10/524,048
March 6, 2008

and the groups Q and W are independently selected from hydrogen or R as is defined above, and Q and W may combine to form a cycle,

the groups Z and Y may combine to form a cycle,

the groups R1 to R5 may not combine together to form a cycle,

with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different,

with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may not be trifluoroacetamido, or acetamido,

with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-amine]benzyl ester (ODmab), N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),

with the further proviso that ~~when the compound is of the 2-deoxy-2-aminoglucose configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [-CH₂-CO₂H] or lactate ether [-CH(CH₃)-CO₂H] or an ester or amide derivative thereof; and with the proviso that the group R may not be or contain another saccharide moiety; and wherein the optional substituents are selected from the group consisting of OH, NO, NO₂, NH₂, N₃, halogen, CF₃, CHF₂, CH₂F, nitrile, alkoxy, aryloxy, amidine, guanidiniums, carboxylic acid, carboxylic acid ester, carboxylic acid amide, aryl, cycloalkyl, heteroalkyl, heteroaryl, aminoalkyl, aminodialkyl, aminotrialkyl, aminoacyl, carbonyl, substituted or unsubstituted imine, sulfate, sulfonamide, phosphate, phosphoramidate, hydrazide, hydroxamate, hydroxamic acid, heteroaryloxy, aminoalkyl, aminoaryl, aminoheteroaryl, thioalkyl, thioaryl or thioheteroaryl, which may be further substituted.~~

2. (Currently Amended) The compound of claim 1 ~~which is a derivative of a pyranose form of a monosaccharide and~~ wherein n is 1.


WEST et al
Appl. No. 10/524,048
March 6, 2008

3. (Cancelled).

4. (Currently Amended) The compound of claim 2, wherein

n is 1,

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;



and the groups Q and W are independently selected from hydrogen or R as is defined above, with the proviso that Y and Z may not both be hydrogen and where two groups in the compound of formula I are N(Y)Z, these groups are different, the groups Z and Y may combine to form a cycle, the groups R1 to R5 may not combine together to form a cycle, with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different,

WEST et al
Appl. No. 10/524,048
March 6, 2008


with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may not be trifluoroacetamido or acetamido,

with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-amino]benzyl ester (ODmab), N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),


with the further proviso that when the scaffold is of the 2-deoxy-2-aminoglucose configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [-CH₂-CO₂H] or lactate ether [-CH(CH₃)-CO₂H] or an ester or amide derivative thereof.

5. (Currently Amended) The compound of any one of claims 1-4 wherein the heteroarylalkyl is substituted by a moiety selected from the group consisting of OH, NO, NO₂, NH₂, N₃, halogen, CF₃, CHF₂, CH₂F, nitrile, alkoxy, aryloxy, amidine, guanidiniums, carboxylic acid, carboxylic acid ester, carboxylic acid amide, aryl, cycloalkyl, heteroalkyl, heteroaryl, aminoalkyl, aminodialkyl, aminotrialkyl, aminoacyl, carbonyl, substituted or unsubstituted imine, sulfate, sulfonamide, phosphate, phosphoramidate, hydrazide, hydroxamate, hydroxamic acid, heteroaryloxy, aminoalkyl, aminoaryl, aminoheteroaryl, thioalkyl, thioaryl or thioheteroaryl, which may be further substituted, with the proviso that the group R may not be or contain a peptide, protein or amino acid.


6. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




7. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:



8. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




9. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




WEST et al
Appl. No. 10/524,048
March 6, 2008


10. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




11. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




12. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




13. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




14. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




15. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




16. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




17. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




18. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:




19. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:



20. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:



21. (Withdrawn) A method of preparing a compound according to claim 1, said method comprising reduction of:



22. (Withdrawn) A method of synthesizing a compound of claim 1 wherein the compound is immobilised to a support.

WEST et al

Appl. No. 10/524,048


March 6, 2008










23. (Withdrawn) The method of claim 22, wherein the compound is immobilised to the support through a hydroxyl group.

24 (Withdrawn) The method of claim 23, wherein the support is selected from the group consisting of derivatised polystyrene, tentagel, wang resin, MBHA resin, aminomethylpolystyrene, rink amide resin, DOX-mpeg and polyethylene glycol.

25. (Previously Presented) The compound of claim 1, wherein R1 is selected from the group consisting of

WEST et al
Appl. No. 10/524,048
March 6, 2008





WEST et al
Appl. No. 10/524,048
March 6, 2008


















26. (Previously Presented) The compound of claim 1, wherein one of the R moieties in OR is selected from the group consisting of


Methyl

Y1

Ethyl


Y2






WEST et al
Appl. No. 10/524,048
March 6, 2008


27. (Currently Amended) The compound of claim 1, wherein Y is selected from the group consisting of




Z1




Z2




Z3




Z4




Z5




Z6




Z7




Z8




Z9




Z10




Z11




Z12




Z13




Z14




Z15



Z16




Z17




Z18


WEST et al
Appl. No. 10/524,048
March 6, 2008




Z19




Z20




Z21




Z22




Z23




Z24




Z25




Z26




Z27




Z28




Z29




Z30



Z31












Z32




Z33


WEST et al
Appl. No. 10/524,048
March 6, 2008

**Z34****Z35****Z36****Z37****Z38****Z39****Z40****Z41****Z42**


WEST et al
Appl. No. 10/524,048
March 6, 2008




Z43




Z44




Z45




Z46




Z47




Z48




Z49




Z50




Z51




Z52



Z53








Z54




Z55

WEST et al.
Appl. No. 10/524,048
March 6, 2008

**Z56****Z57****Z58****Z59****Z60**

28. (New) A compound of formula 2

**formula 2**

R1 is XR wherein,

X is selected from O; S; S=O and SO₂,

WEST et al
Appl. No. 10/524,048
March 6, 2008


R is a moiety selected from the group consisting of: C1 to C9 alkyl, C2 to C15 alkenyl, C2 to C15 alkynyl, C1 to C15 heteroalkyl, C6 to C15 aryl, C6 to C15 heteroaryl, C6 to C15 arylalkyl or C6 to C15 heteroarylalkyl; which moiety is optionally substituted, cyclic or acyclic, branched and/or linear,

the groups R2 to R5 are selected from OH, OR and N(Y)Z such that:

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OH,

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OR, where R is defined above, with the proviso that when two of the groups R2 to R5 are OR, OR is an ether type moiety and the R groups may not both be methyl or unsubstituted benzyl,

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;



WEST et al
Appl. No. 10/524,048
March 6, 2008

and the groups Q and W are independently selected from hydrogen or R as is defined above, and Q and W may combine to form a cycle,
the groups Z and Y may combine to form a cycle,
the groups R1 to R5 may not combine together to form a cycle,

with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different,

with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may not be trifluoroacetamido, acetamido,

with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, *N*-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),

with the proviso that the group R may not be or contain another saccharide moiety; and wherein the optional substituents are selected from the group consisting of OH, NO, NO₂, NH₂, N₃, halogen, CF₃, CHF₂, CH₂F, nitrile, alkoxy, aryloxy, amidine, guanidiniums, carboxylic acid, carboxylic acid ester, carboxylic acid amide, aryl, cycloalkyl, heteroalkyl, heteroaryl, aminoalkyl, aminodialkyl, aminotrialkyl, aminoacyl, carbonyl, substituted or unsubstituted imine, sulfate, sulfonamide, phosphate, phosphoramidate, hydrazide, hydroxamate, hydroxamic acid, heteroaryloxy, aminoalkyl, aminoaryl, aminoheteroaryl, thioalkyl, thioaryl or thioheteroaryl, which may be further substituted.